

THE CURRENT STATUS OF THE VERTEBRATE DIVERSITY IN AL-DALMAJ MARSH, AL-DIWANIYA PROVINCE

Mohammad K. Mohammad
Iraq Natural History Research Center and Museum, University of Baghdad,
Bab Al-Muadham, P.O. Box 59028, Baghdad, Iraq
Email: amarmkm82@yahoo.com

ABSTRACT

A survey conducted at Dalmaj marsh, Al-Diwaniya Province during 2013 revealed that the marsh encounters a considerable part of the Iraqi vertebrate fauna including 147 species belonging to five classes; Pisces, Amphibia, Reptilia, Aves and Mammalia. Some species are of globally conservation importance. The present results are discussed with the pertinent literature.

INTRODUCTION

The global temperature increase may led to changes in the hydroclimatic parameters and have profound impacts on the physical and biological components of the ecosystems in the Euphrates-Tigris Basins as well as on the socio-economic developments of the basin countries (Bozkurt and Sen, 2013). So, studying the biological components of Dalmaj marsh area ecosystems seems necessary to provide a reliable data to compare with during next years which may have more obvious impacts of the global temperature increase. Al-Dalmaj marsh is a large isolated marsh situated at the heart of the Mesopotamian alluvial plain with estimated area of 100000 ha and altitude of less than 20 m (Nature Iraq, 2013), lies to the west of the Tigris River approximately 35 km southwest of Kut City, Wasit Province (Evans, 1994) and about 65 km north east to Al-Diwaniya City, Al-Diwaniya Province. It constitutes an open water lake and marsh with dense reed beds of *Phragmites* and *Typha* in addition to the submerged plants and the plants along the edge of the marsh. It receives water from the Main Outfall Drain (MOD) and discharges again to the MOD with no stable water-level (Salim et al., 2009). The depth of water ranges from less than 0.5 m in the banks of marshy areas to about 1.5-2 m in the center of the lake.

Al-Dalmaj marsh is an important area for the Iraqi biodiversity since it lies within a semi-desert area and comprises terrestrial and aquatic habitats allowing a wide range of biodiversity components. The vertebrates are the most prominent group in the marsh especially birds and fishes and for a lesser extent reptiles and mammals. However, it is still poorly studied and further scientific work is required to understand the biodiversity and the relationships among the biotic and abiotic factors in the area (Nature Iraq, 2013).

The aim of this work is to throw a light on the current situation of vertebrates living in the area, and to provide assessment notes on certain species from the conservation point of view.

MATERIALS AND METHODS

Site description: The following brief description depends mainly on Nature Iraq (2013). Dalmaj wetland is located 40 km east of Diwaniya city and 40 km southwest of Kut city. It

Vertebrate Diversity Dalmaj Marsh

includes both terrestrial habitats ranging from arid areas to true desert with sand dunes, and a large body of water that can be divided into an open-water lake reaching depths exceeding 2 m and true marshes with dense reed beds and shallower water (less than 1 m). Embankments surround the marsh to contain the body of water. The southern section of Dalmaj is mainly mudflats, featuring *Phragmites* and *Typha* reed beds in addition to submerged plants with occasional dry ground scattered with bushes and terrestrial species. The eastern part of the site includes much of the open and deeper Dalmaj Lake that lies within the embankment. To the east of the embankment there are shallow, salty marshes with a dense strip of reed beds and *Tamarix* bushes. The freshwater marshes in the northern part of the site are defined by rich plant cover, such as *Phragmites* and *Typha* reed beds and *Tamarix* in drier areas. These marshes have clear, transparent waters and submerged plants, which provide excellent protection for juvenile fish and offer high oxygen production

Eight field trips (2/season) of 2-3 days each were conducted to Al-Dalmaj marsh lies within the Al-Diwaniya Province during the year 2013 to report on the vertebrate species present in the area. The data presented here is mainly the author's personal observations, direct collection of specimens, photographing, observing species in the field with naked eyes or by the aid of a binocular, examining of remains and traces of vertebrates and interviews with hunters and locals in the area. Specific identifications were possible following Coad (2010) for fishes, Khalaf (1959) for herpetofauna, Salim et al. (2006) for birds, and Harisson (1968, 1981) for mammals.

RESULTS AND DISCUSSION

Table 1 shows that the vertebrate groups in Dalmaj marsh comprise five classes, Class Pisces, Class Amphibia, Class Reptilia, Class Aves and Class Mammalia. This would includes 147 species of vertebrates.

Fishes:

- 1- *Carasobarbus luteus*
- 2- *Barbus xanthopterus*
- 3- *Mesopotamichthys sharpeyi*
- 4- *Cyprinus carpio**
- 5- *Tilapia zilli***
- 6- *Liza abu*
- 7- *Silurus triostegus*

Amphibians:

- 8- *Bufo viridis*
- 9- *Rana ridibunda*

Reptiles:

- 10- *Maurymes caspica*
- 11- *Laudakia persica*
- 12- *Acanthodactylus* sp.
- 13- *Gymnodactylus scaber*
- 14- *Mabuya aurata*
- 15- *Natrix tessellata*
- 16- *Eryx jaculus*

Mohammad K. Mohammad

Birds:

- 17- *Tachybaptus ruficollis*
- 18- *Phalacrocorax carbo*
- 19- *Pelecanus onocrotalus*
- 20- *Botaurus stellaris*
- 21- *Ixobrychus minutus*
- 22- *Nycticorax nycticorax*
- 23- *Egretta garzetta*
- 24- *Ardeola ralloides*
- 25- *Bubulcus ibis*
- 26- *Ardea cinerea*
- 27- *Ardea purpuria*
- 28- *Ciconia ciconia*
- 29- *Plegadis falcinellus*
- 30- *Anser erythrops*
- 31- *Anser anser*
- 32- *Tadorna ferruginea*
- 33- *Tadorna tadotna*
- 34- *Anas penelope*
- 35- *Anas strepera*
- 36- *Anas crecca*
- 37- *Anas platyrhynchos*
- 38- *Anas acuta*
- 39- *Anas clypeata*
- 40- *Marmaronetta angustirostris*
- 41- *Netta rufina*
- 42- *Aythya nyroca*
- 43- *Aythya ferina*
- 44- *Milvus migrans*
- 45- *Neophron percnopterus*
- 46- *Circus aeruginosus*
- 47- *Buteo rufinus*
- 48- *Aquila clanga*
- 49- *Falco tinnunculus*
- 50- *Francolinus francolinus*
- 51- *Rallus aquaticus*
- 52- *Gallinula chloropus*
- 53- *Fulica atra*
- 54- *Porphyrio porphyrio*
- 55- *Grus grus*
- 56- *Chlamydotis undulata*
- 57- *Himantopus himantopus*
- 58- *Recurvirostra avosetta*
- 59- *Glareola pratincola*
- 60- *Charadrius dubius*
- 61- *Charadrius alexandrinus*
- 62- *Hoplopterus spinosus*
- 63- *Hoplopterus indicus*
- 64- *Chettusia leucura*
- 65- *Calidris minuta*
- 66- *Calidris temminckii*

Vertebrate Diversity Dalmaj Marsh

- 67- *Calidris alpina*
- 68- *Gallinago gallinago*
- 69- *Limosa limosa*
- 70- *Tringa totanus*
- 71- *Tringa stagnatilis*
- 72- *Tringa nebularia*
- 73- *Tringa ochropus*
- 74- *Tringa glareola*
- 75- *Larus ridibundus*
- 76- *Larus genei*
- 77- *Larus armenicus*
- 78- *Sterna caspia*
- 79- *Sterna hirundo*
- 80- *Chlidonia hybridus*
- 81- *Chlidonia leucopterus*
- 82- *Pterocles alchata*
- 83- *Columba livia*
- 84- *Columba palumbus*
- 85- *Streptopelia decaocto*
- 86- *Streptopelia turtur*
- 87- *Streptopelia senegalensis*
- 88- *Athene noctua*
- 89- *Tyto alba*
- 90- *Caprimulgus aegyptius*
- 91- *Halcyon smyrnensis*
- 92- *Ceryle rudis*
- 93- *Alcedo atthis*
- 94- *Merops superciliosus*
- 95- *Coracias benghalensis*
- 96- *Upupa epops*
- 97- *Ammomanes deserti*
- 98- *Galerida cristata*
- 99- *Calandrella brachydactyla*
- 100- *Riparia riparia*
- 101- *Hirundo rustica*
- 102- *Anthus trivialis*
- 103- *Anthus spinoletta*
- 104- *Motacilla flava*
- 105- *Motacilla alba*
- 106- *Pycnonotus leucogenys*
- 107- *Hypocolius ampelinus*
- 108- *Erithacus rubicula*
- 109- *Saxicola rubetra*
- 110- *Saxicola torquata*
- 111- *Oenanthe deserti*
- 112- *Cisticola juncidis*
- 113- *Prinia gracilis*
- 114- *Acrocephalus griseldis*
- 115- *Hippolias pallida*
- 116- *Sylvia mystacea*
- 117- *Phylloscopus collybita*

Mohammad K. Mohammad

- 118- *Turdoides altirostris*
- 119- *Turdoides caudatus*
- 120- *Lanius collurio*
- 121- *Lanius nubicus*
- 122- *Corvus frugilegus*
- 123- *Corvus corone*
- 124- *Sturnus vulgaris*
- 125- *Passer domesticus*
- 126- *Passer hispaniolensis*
- 127- *Passer moabiticus*.
- 128- *Carduelis carduelis*
- 129- *Carduelis chloris*
- 130- *Rhodospiza obsoleta*
- 131- *Emberiza hortulana*
- 132- *Emberiza schoeniclus*
- 133- *Miliaria calandra*.

Mammals:

- 134- *Pipistrellus kuhlii*
 - 135- *Taphozous nudiventris*
 - 136- *Hystrix indica*
 - 137- *Mus musculus*
 - 138- *Rattus rattus*
 - 139- *Rattus norvegicus**
 - 140- *Lepus capensis*
 - 141- *Canis aureus*
 - 142- *Canis lupus*
 - 143- *Vulpes vulpes*
 - 144- *Hyaena hyaena*
 - 145- *Herpestes auropunctatus*
 - 146- *Meles meles*
 - 147- *Sus scrofa*
-

*exotic species

**invasive species

Dalmaj is rich in biodiversity, being a wintering ground for numerous waterfowl and a main breeding area for Marbled Duck *Marmaronetta angustirostris*, Ferruginous Duck *Aythya nyroca*, and Red-crested Pochard *Netta rufina*, three of the four known breeding ducks in Iraq, and a major breeding site for the Endangered Basra Reed Warbler (nature Iraq, 2013). The list of species included in table 1 indicates that the marsh contains a considerable part of the Iraqi avifauna which counts to more than 400 species. The number of the recorded birds in this study is 117 while Nature Iraq (2013) mentioned that 140 migratory and resident birds were seen in the marsh. In general, it seems that the biodiversity components of the marsh is rather in a good condition. Some species in table 1 are of certain conservation importance including *Tilapia zilli* (Gervais, 1848), *Marmaronetta angustirostris* (Menetries, 1823) L. Reichenbach, 1853, *Aythya nyroca* and *Acrocephalus griseldus*.

Tilapia zilli: (fig. 1), this is an invasive fish. It is considered a potential competitor with native fish for food and spawning areas (Molnar, 2008). In Dalmaj marsh, the author was able to observe it is widely distributed along the marsh shore. On 28.5.2013 a couple of male and

Vertebrate Diversity Dalmaj Marsh

female fishes were surrounded with a 30 cm diameter ball of hundreds of apparently newly hatched fishes just 1.5-2 m off the shore. The author counted 7 pairs of parents with their juvenile balls in about 50 m distance of the shore. The parents ran away only a little when thrown with a stone and the juvenile ball tends to decrease the diameter and become denser, but sooner the parents return back to their juveniles defending them.



Fig. 1: *Tilapia zilli* from al-Dalmaj marsh.

***Marmaronetta angustirostris*:** It is a resident bird in suitable habitats in the middle and south and for a lesser extent in the north, and breeds from May to July (Allouse, 1960). IUCN red list for the year 2012 considered this duck globally vulnerable (A2cd+3cd+4cd ver 3.1) with decreasing population trend (Salim *et al.*, 2009; BirdLife international, 2012). The marsh proved to be a suitable breeding site for this species and the author was able to take pictures of duckling with parents (fig. 2). However, severe hunting practices is going on in the area and thousands of this duck was brought to local markets during the period from September to March (fig. 3). The hunting pressure is extremely critical during September to end of November at which hunting was practically targeted this duck since it the only available game bird in reasonable number.

Mohammad K. Mohammad



Fig.2: Mother marbled teal *Marmaronetta angustirostris* and eight ducklings in Dalmaj marsh



Fig. 3: Marbled teals brought from Dalmaj marsh sold at a local market in Baghdad city.

***Aythya nyroca*:** (fig. 4) It is considered globally Near Threatened ver 3.1 with a decreasing population trend (IUCN, 2013). The author's personal observations indicated that this duck breeds in Al-Attariya marsh southeast of Baghdad not far from Dalmaj marsh since the 70s decade of the last century. George and Vieliard (1970) recorded only 31 individuals, then Scott and Carp (1982) could not find this species in their survey to Iraqi marshes. It was proved later that it breeds in some other sites in the middle and south of Iraq including Dalmaj

Vertebrate Diversity Dalmaj Marsh

marsh and southern Iraqi marshes. Howeverm Salim *et al.* (2009) stated that further surveys will show this duck to be more common and more widespread in the Mesopotamian marshes. Fig. (5) shows hundreds of ferruginous ducks captured in Al-Dalmaj marsh and sold at a local market in Baghdad city.



Fig 4: Male Ferruginous duck *Aythya nyroca*



Fig. 5: Hundreds of ferruginous ducks brought from Dalmaj marsh sold at a local market in Baghdad

Mohammad K. Mohammad

Acrocephalus griseldis: The Basra Reed Warbler is a globally endangered bird (IUCN, 2013). Iraq is known to encounter more than 90% of the world population of the bird (Richardson and Hussain, 2006). Nature Iraq (2013) counted up to 900 breeding pairs in Dalmaj marsh. Many nests were noted by the author among reed beds in the marsh.

LITERATURE CITED

- Bozkurt, D. and Sen, O. L. 2013. Climate change impacts in the Euphrates–Tigris Basin based on different model and scenario simulations. *Journal of Hydrology*, 480: 149–161.
- Coad, B. W. 2010 *Freshwater fishes of Iraq*. Pensoft. Sofia-Moscow, 274 pp.
- Evans, M. I. 1994. *Important Bird Areas in the Middle East*. Cambridge: BirdLife International.
- George P. V. and Vielliard J. 1970. *Midwinter Observations on Birds of Central and South Iraq*. 1968. Baghdad, Iraq: Iraq Natural History Museum Bull. Vol. IV (4).
- Harrison, D. L. 1968 *The mammals of Arabia*. Ernest Benn Ltd., London, 3vols.
- Harrison, D. L. 1981 *The mammals of the Arabian Gulf*. George Allen and Unwin, London, 92pp.
- IUCN 2013. *IUCN Red List of Threatened Species*. Version 2013.2. <www.iucnredlist.org>.
- Khalaf, K. T. 1959 *Reptiles of Iraq with some notes on the amphibians*. ArRabitta Press, Baghdad, 96 pp.
- Molnar, J. L., Gamboa, R. L., Revenga, C. and Spalding, M. D. 2008. Assessing the global threat of invasive species to marine biodiversity. *Frontiers in Ecology and the Environment* 6: 485–492.
- Nature Iraq 2013. available online:
http://www.natureiraq.org/uploads/9/2/7/0/9270858/dalmaj_me10_22_maanna.pdf.
- Richardson, C.J., and N.A. Hussain. 2006. Restoring the Garden of Eden: An ecological assessment of the marshes of Iraq. *BioScience* 56 (6):477-489.
- Salim, M., Porter, R. and Rubec, C. 2009. A summary of birds recorded in the marshes of southern Iraq, 2005–2008. *BioRisk* 3: 205–219. doi: 10.3897/biorisk.3.14. www.pensoftonline.net/biorisk.
- Salim, M. A., Porter, R. F., Christensen, S. Schiermacker-Hansen, P. and Al-Jbour, S. 2006. *Field Guide to the birds of Iraq*. Amman: Nature Iraq & BirdLife International. (In Arabic).
- Scott, D. A. and Carp, E. 1982. A midwinter survey of wetlands in Mesopotamia, Iraq: 1979. *Sandgrouse*, No. 4.

الواقع الحالي لتنوع الفقريات في هور الدلمج، محافظة الديوانية

محمد كاظم محمد

مركز بحوث ومتحف التاريخ الطبيعي، جامعة بغداد، باب المعظم، ص. ب. ٥٩٠٢٨،
بغداد، العراق

Email: amarmkm82@yahoo.com

الخلاصة

بين مسح اجري في هور الدلمج محافظة الديوانية خلال عام ٢٠١٣ بان هذا الهور يشتمل على جزء مهم من المجموعة الحيوانية الفقرية العراقية ويضم ١٤٧ نوعا تعود الى خمسة اصناف هي الاسماك والبرمائيات والزواحف والطيور واللبائن. تبين ان بعض الانواع المسجلة ذات اهمية من ناحية الاهمية العالمية للمحافظة. نوقشت النتائج الحالية في ضوء البحوث ذات العلاقة.